

Rank-width and Well-quasi-ordering of Symmetric or Skew-symmetric Matrices

Sang-il Oum

Program in Applied and Computational Mathematics
Princeton University

June 16, 2005

Séminaire de Combinatoire Algébrique et Géométrie

Well-quasi-ordering

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

A set Q is **well-quasi-ordered** by an ordering \preceq if
 \forall infinite sequences a_1, a_2, \dots of Q ,

$\exists i, j$ such that

$$i < j \text{ and } a_i \preceq a_j.$$

Equivalently, we say: \leq is an **well-quasi-ordering** of Q .

Examples:

- 1 {positive integers} is well-quasi-ordered by \leq .
- 2 {integers} is NOT well-quasi-ordered by \leq .

Overview

common generalization { symmetric or skew-symmetric matrices,
delta-matroids representable over a finite field.

(1) Robertson and Seymour (1990)

Graphs + minors

Tree-width and well-quasi-ordering

(2) Geelen, Gerards, and Whittle (2002)

Matroids representable over a finite field + minors

Branch-width and well-quasi-ordering

(3) Oum (2005)

Graphs + Vertex-minors

Rank-width (or Clique-width) and well-quasi-ordering

(3) implies (2) for $GF(2)$. (2) or (3) implies (1).

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Preview of main tools

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing



Ph.D. thesis: An Algebraic Theory of Graphs

*Develop the theory of **Chain-groups**
(unaware of matroids at that time)*

A matroid is **representable**

\Leftrightarrow There is a **chain-group** representing the matroid.

Delta-matroids: (sort of) relaxation of matroids.

A delta-matroid is **representable** (by def) if

*it is equivalent to a delta-matroid represented by
some symmetric or skew-symmetric matrices.*

We develop the theory of **Lagrangian Chain-groups**.

A delta-matroid is **representable**

\Leftrightarrow There is a **Lagrangian chain-group** representing it.

Preliminaries

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

- Transpose of a matrix: A^t .
- **Symmetric** matrices: $A = A^t$.
- **Skew-symmetric** matrices:
 - 1 $A = -A^t$.
 - 2 0 on the diagonal entries.
(We require this for matrices over $\text{GF}(2)$.)
- Matrices of **symmetric type**: either symmetric or skew-symmetric
- Submatrix of A : $A[X, Y]$ (rows in X , columns in Y)
- **Principal Submatrix** of A : $A[X] = A[X, X]$

(Principal) Pivoting

Let A : $V \times V$ matrix.

If $A[X]$ is nonsingular, then

$$A = \begin{matrix} & X & Y \\ X & \alpha & \beta \\ Y & \gamma & \delta \end{matrix} \implies A * X = \begin{matrix} & X & Y \\ X & \alpha^{-1} & \alpha^{-1}\beta \\ Y & -\gamma\alpha^{-1} & \delta - \gamma\alpha^{-1}\beta \end{matrix}$$

- 1 If A is skew-symmetric, then $A * X$ is skew-symmetric.
- 2 If A is symmetric, we can obtain a symmetric matrix by negating columns of X in $A * X$.

Pivot-minors of a skew-symmetric matrix A : $A * X[Y]$

Pivot-minors of a symmetric matrix A : Symmetric matrices obtained by resigning of columns in $A * X[Y]$.

Rank-width

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Let $M: V \times V$ matrices of symmetric type

- **Rank-decomposition**: a pair (T, \mathcal{L}) of
 - 1 a subcubic tree T
 - 2 a bijection $\mathcal{L} : V \rightarrow \{\text{leaves of } T\}$.
- **Width** of an edge e of T : $\text{rank } M[\mathcal{L}^{-1}(A_e), \mathcal{L}^{-1}(B_e)]$
 (A_e, B_e) : partition of leaves of T induced by $T \setminus e$.
- **Width** of a rank-decomposition (T, \mathcal{L}) :
Maximum width of edges of T .
- **Rank-width** of M :
Minimum width of all its rank-decompositions

Our theorem (simplified, matrix version)

- $A \times B$ Matrix over F : a mapping from $A \times B$ to F .
- **Isomorphism**
from a $V \times V$ matrix M to a $W \times W$ matrix N :
a bijection μ from V to W such that
 $M(x, y) = N(\mu(x), \mu(y))$.

Let k : constant, F : **finite** field.

Matrices over F of symmetric types having rank-width $\leq k$
are **well-quasi-ordered** by pivot-minors.

More precisely, if M_1, M_2, \dots are matrices of symmetric type
over F having rank-width $\leq k$,
then there exist i and j such that
 $i < j$ and M_i is **isomorphic to a pivot-minor** of M_j .

Chain-groups

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Let V : finite set, F : field.

Let $K = F^2 = \left\{ \begin{pmatrix} a \\ b \end{pmatrix} : a, b \in F \right\}$: 2-dim vector space over F .

- **Chain**: function from V to K .
- Sum of chains: $f + g$
- Scalar product cf when $c \in F$.
- **Chain-group**: set of chains on V to K closed under sum and scalar product. (subspace of K^V)

Note:

Tutte's chain: function from V to F .

Lagrangian chain-groups

We give a bilinear form to chains.

- Let $\langle \begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} c \\ d \end{pmatrix} \rangle_K = ad + bc$ (symmetric)
or $\langle \begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} c \\ d \end{pmatrix} \rangle_K = ad - bc$. (skew-symmetric)
- For chains f, g on V to K ,

$$\langle f, g \rangle = \sum_{v \in V} \langle f(v), g(v) \rangle_K.$$

- A chain-group N on V to K is called **isotropic**:

$$\forall f, g \in N, \quad \langle f, g \rangle = 0.$$

- A chain-group N on V to K is called **Lagrangian**:
isotropic and $\dim(N) = |V|$.

Idea from both
Chain-groups (Tutte) + Isotropic systems (Bouchet)

Minors of chain-groups

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

For a chain f on V to K ,

Restriction $f \cdot T$: chain on T to K such that

$$f \cdot T(x) = f(x) \text{ for all } x \in T.$$

For a chain-group N on V to K ,

- **Deletion $N \ominus T$:**

$$\{f \cdot (V \setminus T) : f \in N, \langle f(x), \binom{1}{0} \rangle_K = 0 \forall x \in T\}.$$

- **Contraction $N \oslash T$:**

$$\{f \cdot (V \setminus T) : f \in N, \langle f(x), \binom{0}{1} \rangle_K = 0 \forall x \in T\}.$$

Minors of a chain-group N on V to K :

Chain-groups of the form $N \ominus X \oslash Y$.

Properties of Minors of chain-groups

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

- 1 A minor of a minor of N is a minor of N .
- 2 A minor of an **isotropic** chain-group is **isotropic**.
- 3 A minor of a **Lagrangian** chain-group is **Lagrangian**.

Minor-closed classes of chain-groups.

Chain-groups

Isotropic Chain-groups

Lagrangian Chain-groups

Note: Bouchet 1987 introduced isotropic chain-groups.

Algebraic Duality

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Let N : chain-group.

Let $N \cdot T = \{f \cdot T : f \in N\}$,

$N \times T = \{f \cdot T : f \in N, f(x) = 0 \forall x \notin T\}$,

$N^\perp = \{g : \langle f, g \rangle = 0 \forall f \in N\}$.

Theorem (Same proof by Tutte works)

$$(N \cdot T)^\perp = N^\perp \times T.$$

$$(N \otimes T)^\perp = N^\perp \otimes T,$$

$$(N \oplus T)^\perp = N^\perp \oplus T.$$

Connectivity

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

For a chain-group N on V to K ,

$$\lambda_N(U) = \frac{\dim N - \dim(N \times (V \setminus U)) - \dim(N \times U)}{2}$$

- **Symmetric:** $\lambda_N(X) = \lambda_N(V \setminus X)$.
- **Submodular:** $\lambda_N(X) + \lambda_N(Y) \geq \lambda_N(X \cap Y) + \lambda_N(X \cup Y)$.

If N is Lagrangian, then

$$\lambda_N(U) = |V| - \dim(N \times U).$$

Branch-width of chain-groups

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

For a chain-group on V to K ,

- **Branch-decomposition**: a pair (T, \mathcal{L}) of
 - 1 a subcubic tree T
 - 2 a bijection $\mathcal{L} : V \rightarrow \{\text{leaves of } T\}$.
- **Width** of an edge e of T : $\lambda_N(\mathcal{L}^{-1}(A_e))$.
 (A_e, B_e) : partition of leaves of T induced by $T \setminus e$.
- **Width** of a branch-decomposition (T, \mathcal{L}) :
Maximum width of edges of T .
- **Branch-width** of M :
Minimum width of all its branch-decompositions

If M is a minor of N , then

$$(\text{branch-width of } M) \leq (\text{branch-width of } N).$$

Our theorem (simplified)

- **Simple Isomorphism**

from a chain-group N on V to K to a chain-group M on W to K : a bijection μ from V to W such that $N = \{f \circ \mu : f \in M\}$.

Let k : constant, F : **finite** field. $K = F^2$.

Lagrangian chain-groups over F having branch-width $\leq k$ are **well-quasi-ordered** by minors.

More precisely, if N_1, N_2, \dots are Lagrangian chain-groups over F having branch-width $\leq k$, then there exist i and j such that $i < j$ and N_i is **simply isomorphic to a minor** of N_j .

Delta-matroids

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Axiom for bases of matroids (E, \mathcal{B})

- for B_1 and $B_2 \in \mathcal{B}$,
if $x \in B_1 \setminus B_2$, then
 $\exists y \in B_2 \setminus B_1$ such that $B_1 \Delta \{x, y\} \in \mathcal{B}$
- $\mathcal{B} \neq \emptyset$.

Axiom for **feasible** sets of **delta-matroids** (V, \mathcal{F})

- for F_1 and $F_2 \in \mathcal{F}$,
if $x \in F_1 \Delta F_2$, then
 $\exists y \in F_1 \Delta F_2$ such that $F_1 \Delta \{x, y\} \in \mathcal{F}$
- $\mathcal{F} \neq \emptyset$.

Twisting, Deletion, and Minors

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

If $\mathcal{M} = (V, \mathcal{F})$ is a delta-matroid, then $\mathcal{M}' = (V, \{F\Delta X : F \in \mathcal{F}\})$ is also a delta-matroid. This operation is called a **twisting**. We write $\mathcal{M}' = \mathcal{M}\Delta X$.

Two delta-matroids are **equivalent** if one is obtained from another by twisting.

If there is a feasible set Y such that $Y \cap X = \emptyset$, then $\mathcal{M} \setminus X = (V, \{F \in \mathcal{F} : F \cap X = \emptyset\})$ is a delta-matroid. This operation is called a **deletion**.

A delta-matroid obtained by twisting and deletion from \mathcal{M} is called a **minor** of \mathcal{M} .

Representable Delta-matroids

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

For a $V \times V$ matrix A of symmetric type,
Let $\mathcal{F} = \{X : A[X] \text{ is nonsingular}\}$.

Theorem (Bouchet, 1987)

$\mathcal{M}(A) = (V, \mathcal{F})$ is a delta-matroid.

Representable delta-matroids:

Delta-Matroids of the form $\mathcal{M}(A)\Delta X$

His proof:

Matrices of symmetric type

⇒ **Isotropic chain-groups**

⇒ Delta-matroids

Matrices to Chain-groups

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Let $A = (a_{ij})_{i,j \in V}$ be a matrix of symmetric type over F .

Two chains a, b on V are called **supplementary** if
 $\langle a(v), b(v) \rangle_K = \langle a(v), a(v) \rangle_K = \langle b(v), b(v) \rangle_K = 0$.

$$\langle \cdot, \cdot \rangle_K = \begin{cases} \text{symmetric} & \text{if } A \text{ is skew-symmetric,} \\ \text{skew-symmetric} & \text{if } A \text{ is symmetric.} \end{cases}$$

Let f_i : a chain on V to K such that

$$f_i(j) = \begin{cases} a_{ij} \begin{pmatrix} 1 \\ 0 \end{pmatrix} & \text{if } i \neq j, \\ a_{ij} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \end{pmatrix} & \text{if } i = j. \end{cases}$$

Let N : a chain-group **spanned by** $\{f_i : i \in V\}$.

If $i \neq j$, then $\langle f_i, f_j \rangle = a_{ij} \langle \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \rangle_K + a_{ji} \langle \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix} \rangle_K = 0$.

If $i = j$, then $\langle f_i, f_i \rangle = 0$. **N is isotropic.**

Since $\dim(N) = |V|$, **N is Lagrangian.**

Matrices to Chain-groups

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Let $A = (a_{ij})_{i,j \in V}$ be a matrix of symmetric type over F .
Two chains a, b on V are called **supplementary** if
 $\langle a(v), b(v) \rangle_K = \langle a(v), a(v) \rangle_K = \langle b(v), b(v) \rangle_K = 0$.

$$\langle \cdot, \cdot \rangle_K = \begin{cases} \text{symmetric} & \text{if } A \text{ is skew-symmetric,} \\ \text{skew-symmetric} & \text{if } A \text{ is symmetric.} \end{cases}$$

Let f_i : a chain on V to K such that

$$f_i(j) = \begin{cases} a_{ij} \begin{pmatrix} 1 \\ 0 \end{pmatrix} & \text{if } i \neq j, \\ a_{ij} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \end{pmatrix} & \text{if } i = j. \end{cases}$$

Let N : a chain-group **spanned by** $\{f_i : i \in V\}$.

If $i \neq j$, then $\langle f_i, f_j \rangle = a_{ij} \langle \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \rangle_K + a_{ji} \langle \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix} \rangle_K = 0$.

If $i = j$, then $\langle f_i, f_i \rangle = 0$. **N is isotropic.**

Since $\dim(N) = |V|$, **N is Lagrangian.**

Matrices to Chain-groups

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Let $A = (a_{ij})_{i,j \in V}$ be a matrix of symmetric type over F .
Two chains a, b on V are called **supplementary** if
 $\langle a(v), b(v) \rangle_K = \langle a(v), a(v) \rangle_K = \langle b(v), b(v) \rangle_K = 0$.

$$\langle \cdot, \cdot \rangle_K = \begin{cases} \text{symmetric} & \text{if } A \text{ is skew-symmetric,} \\ \text{skew-symmetric} & \text{if } A \text{ is symmetric.} \end{cases}$$

Let f_i : a chain on V to K such that

$$f_i(j) = \begin{cases} a_{ij}a(v) & \text{if } i \neq j, \\ a_{ij}a(v) + b(v) & \text{if } i = j. \end{cases}$$

Let N : a chain-group **spanned by** $\{f_i : i \in V\}$.

If $i \neq j$, then $\langle f_i, f_j \rangle = a_{ij} \langle \binom{1}{0}, \binom{0}{1} \rangle_K + a_{ji} \langle \binom{0}{1}, \binom{1}{0} \rangle_K = 0$.

If $i = j$, then $\langle f_i, f_i \rangle = 0$. **N is isotropic.**

Since $\dim(N) = |V|$, **N is Lagrangian.**

Isotropic Chain-groups to Delta-matroids

Let N : isotropic chain-groups on V to K .

Theorem (Bouchet)

Say $F \in \mathcal{F}$ if $f \equiv 0$ is the only chain in N satisfying

- $\langle f(x), \begin{pmatrix} 1 \\ 0 \end{pmatrix} \rangle_K = 0$ for all $x \notin F$,
- $\langle f(x), \begin{pmatrix} 0 \\ 1 \end{pmatrix} \rangle_K = 0$ for all $x \in F$.

(V, \mathcal{F}) is a delta-matroid.

Moreover, if N is from a matrix A of symmetric type, F is feasible if and only if $A[X]$ is nonsingular.

Matrices of symmetric type

⇒ Lagrangian chain-groups

⇒ Isotropic chain-groups

⇒ Delta-matroids

Reverse
direction?

Isotropic Chain-groups to Delta-matroids

Let N : isotropic chain-groups on V to K .

Theorem (Bouchet)

Say $F \in \mathcal{F}$ if $f \equiv 0$ is the only chain in N satisfying

- $\langle f(x), a(v) \rangle_K = 0$ for all $x \notin F$,
- $\langle f(x), b(v) \rangle_K = 0$ for all $x \in F$.

(V, \mathcal{F}) is a delta-matroid.

Moreover, if N is from a matrix A of symmetric type, F is feasible if and only if $A[X]$ is nonsingular.

Matrices of symmetric type

⇒ Lagrangian chain-groups

⇒ Isotropic chain-groups

⇒ Delta-matroids

Reverse
direction?

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Isotropic Chain-groups to Delta-matroids

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Let N : isotropic chain-groups on V to K .

Theorem (Bouchet)

Say $F \in \mathcal{F}$ if $f \equiv 0$ is the only chain in N satisfying

- $\langle f(x), a(v) \rangle_K = 0$ for all $x \notin F$,
- $\langle f(x), b(v) \rangle_K = 0$ for all $x \in F$.

(V, \mathcal{F}) is a delta-matroid.

Moreover, if N is from a matrix A of symmetric type, F is feasible if and only if $A[X]$ is nonsingular.

Matrices of symmetric type

⇒ Lagrangian chain-groups

⇒ Isotropic chain-groups

⇒ Delta-matroids

Reverse
direction?

Isotropic Chain-groups to Delta-matroids

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Let N : isotropic chain-groups on V to K .

Theorem (Bouchet)

Say $F \in \mathcal{F}$ if $f \equiv 0$ is the only chain in N satisfying

- $\langle f(x), a(v) \rangle_K = 0$ for all $x \notin F$,
- $\langle f(x), b(v) \rangle_K = 0$ for all $x \in F$.

(V, \mathcal{F}) is a delta-matroid.

Moreover, if N is from a matrix A of symmetric type, F is feasible if and only if $A[X]$ is nonsingular.

Matrices of symmetric type

⇒ Lagrangian chain-groups

⇒ Isotropic chain-groups

⇒ Delta-matroids

Reverse
direction?

Twisting in Isotropic Chain-groups

Isotropic chain-group N \longrightarrow Delta-matroid \mathcal{M}

\parallel

twisting \downarrow

Same isotropic chain-group \longleftarrow Delta-matroid $\mathcal{M}\Delta X$.

If \mathcal{M} : represented by N with supplementary chains a and b , then $\mathcal{M}\Delta X$ is represented by N with a' , b' .

	$x \notin X$	$x \in X$	$+b(x)$ if \langle , \rangle_K is symmetric, $-b(x)$ if \langle , \rangle_K is skew-symmetric.
$a'(x)$	$a(x)$	$\pm b(x)$	
$b'(x)$	$b(x)$	$a(x)$	

Representable delta-matroids \Rightarrow Lagrangian chain-groups

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Lagrangian Chain-groups to Matrices

Lagrangian chain-groups to Representable delta-matroids

For a Lagrangian chain-group N ,

- Choose supplementary chains a, b so that \emptyset is feasible in the delta-matroid of N (by twisting).
- For each $v \in V$, there exists a unique chain $f_v \in N$ such that

$$\langle a(v), f_v(w) \rangle_K = \begin{cases} 0 & \text{if } v \neq w \\ 1 & \text{if } v = w. \end{cases}$$

- Construct a $V \times V$ matrix $A = (\langle f_i(j), b(j) \rangle_K : i, j \in V)$.

Then the matrix with a, b represents N .

Matrices of symmetric type

\Leftrightarrow Lagrangian chain-groups

\Leftrightarrow Representable Delta-matroids

Connectivity

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

If N : Lagrangian chain-group from a matrix A , then

$$\text{rank } A[X, V \setminus X] = \lambda_N(X).$$

Reminder: $\lambda_N(X) = |X| - \dim(N \times X)$ for Lagrangian chain-group N .

Corollary

*Rank-width of a matrix of symmetric type
= Branch-width of a Lagrangian chain-group from the matrix*

Minors

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Theorem

For an isotropic chain-group N on V to K and $v \in V$, if

- a, b : supplementary chains s.t. $a(v) \in \left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \end{pmatrix} \right\}$,
- \mathcal{M} : delta-matroid represented by N with a, b ,

then $N \circledast \{v\}$ with $a \cdot (V \setminus \{v\}), b \cdot (V \setminus \{v\})$ represents one of the following delta-matroids.

- $\mathcal{M} \setminus \{v\}$ if there is a feasible set not containing v ,
- $\mathcal{M} \Delta \{v\} \setminus \{v\}$ otherwise.

$N \circledast \{v\}$ with $a \cdot (V \setminus \{v\}), b \cdot (V \setminus \{v\})$ represents one of the following delta-matroids.

- $\mathcal{M} \Delta \{v\} \setminus \{v\}$ if there is a feasible set containing v ,
- $\mathcal{M} \setminus \{v\}$ otherwise.

Delta-matroid minors \Leftrightarrow Isotropic Chain-group minors

Pivoting and Twisting

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices


Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing



If N : represented by A with supplementary chains a and b ,
then N is represented by $A * X$ with a' , b' .

	$x \notin X$	$x \in X$	
$a'(x)$	$a(x)$	$\pm b(x)$	$+b(x)$ if \langle , \rangle_K is symmetric, $-b(x)$ if \langle , \rangle_K is skew-symmetric.
$b'(x)$	$b(x)$	$a(x)$	

Pivot-minors \Leftrightarrow Minors of Lagrangian Chain-groups

Our theorem

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

For a well-quasi-ordered set Q with an ordering \preceq ,

- **Q-labeling**: a function from an element set to Q .
- **Q-labeled** chain-group on V : a chain-group with a Q -labeling.
- **Q-minor** of a Q -labeled chain-group N is a minor on V' such that
(Q -labeling of minor)(x) \preceq (Q -labeling of N)(x)
for all $x \in V'$.

If N_1, N_2, \dots are **Q-labeled** Lagrangian chain-groups over F having branch-width $\leq k$,
then there exist i and j such that
 $i < j$ and N_i is **simply isomorphic to a Q-minor** of N_j .

Proof Sketch (Vaguely)

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

- Tutte's linking theorem for Lagrangian chain-groups
- Existence of linked branch-decompositions
Thomas (tree-decomposition), Geelen et al.
(branch-decomposition).
- **Boundary** of an isotropic chain-group N : an ordered basis of N^\perp/N .
- **Boundaried isotropic chain-group**: An isotropic chain-group with a boundary.
- **Sum** of two boundaried isotropic chain-group.
- **Connection types**: describe the sum of two boundaried isotropic chain-group **uniquely**.
- Lemma on Trees by Robertson and Seymour.

(Vaguely) If branch-width is bounded, number of distinct connection types is finite. Using that, show that there is no “minimal” antichain.

Matroids Representable over a Finite Field

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

If \mathcal{M} is a representable matroid, let B be a basis. There is a standard representation:

$$\begin{array}{cc} B & E(\mathcal{M}) \setminus B \\ \left(\begin{array}{cc} 1 & \\ & \ddots & \\ & & 1 & \\ & & & A \end{array} \right) \end{array}$$

Take the following skew-symmetric matrix.

$$S = \begin{array}{cc} B & E(\mathcal{M}) \setminus B \\ \left(\begin{array}{cc} 0 & A \\ -A^t & 0 \end{array} \right) \end{array}$$

Theorem (Bouchet)

$\mathcal{M}(S) \Delta B$ is a delta-matroid whose feasible sets are bases of \mathcal{M} .

Moreover,
Matroid Branch-width
= (Rank-width of S) + 1.


Matroid Minors

⇒ Pivot-minors

⇔ Delta-matroid minors

Representable Matroids of Bounded Branch-width

Let $Q = \{x, y\}$ and $x \not\preceq y$ and $y \not\preceq x$. We assign x for elements in the chosen base, y for other elements.



As a corollary, we conclude that:

Matroids representable over a fixed finite field having bounded branch-width are well-quasi-ordered by matroid minors.

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

Implications

- Matroids (Geelen, Gerards, and Whittle. 2002)

Matroids representable over a fixed finite field having bounded branch-width are well-quasi-ordered by matroid minors.

- Graphs (Robertson and Seymour. 1990)

Graphs representable over a fixed finite field having bounded branch-width are well-quasi-ordered by matroid minors.

- Graphs with Rank-width. (Oum. 2005)

Rank-width of a graph: Rank-width of its adjacency matrix over $GF(2)$.

Pivot-minor of a graph: Pivot-minor of the adjacency matrix.

Open problems — Branch-width, Well-quasi-ordering

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

- Suitable Connectivity functions for Representable Delta-matroids.

My definition of branch-width of delta-matroids representable over F :

Minimum branch-width of Lagrangian chain-groups over F representing the delta-matroid.

Then, F -representable delta-matroids of bounded branch-width are well-quasi-ordered.

- Are delta-matroids representable over a finite field well-quasi-ordered?

This will imply the Graph Minor Theorem.

Open problems — isotropic chain-groups

Rank-width
and WQO

Sang-il Oum

Introduction

Matrices

Chain-groups

Delta-
matroids

Main Theorem

Applications

Closing

- Characterization of Delta-matroids obtained from isotropic chain-groups?

- They are minor-closed.
 - They are either representable delta-matroids or something else(?).
-
- Are isotropic chain-groups of bounded branch-width well-quasi-ordered?
 - Is Tutte's linking theorem true for isotropic chain-groups?

Thank you for your attention!